arXiv:1302.2771v1 [quant-ph] 12 Feb 2013

# Evolution of quasi-Bell states in a qubit-oscillator system with moderately strong coupling

R. Chakrabarti and B. Virgin Jenisha

*Department of Theoretical Physics,*
*University of Madras,*
*Maraimalai Campus, Guindy,*
*Chennai 600 025, India*

## Abstract

We study evolution of bipartite entangled quasi-Bell states in a qubit-oscillator system in the moderately strong coupling regime and also in the presence of a static bias. Using the adiabatic approximation that relies on the separation of the time scales of the two components, the reduced density matrices for the qubit and the oscillator degrees of freedom are obtained in closed forms comprising of linear combinations of Jacobi theta functions. The analytical results based on theta functions are found to be in good agreement with their series counterparts in the said parametric regime. The entropy of the system quantifying the entanglement is computed via the qubit reduced density matrix. The time dependence of this density matrix owes its origin to the bias term. The reduced density matrix of the oscillator is employed for obtaining the phase space distributions such as the Husimi $Q$-function,which, in turn, is utilized for extracting expectation values of antinormally ordered operators.

PACS Number(s): 42.50.-p, 42.50.Pq, 03.65.Ud, 02.30.Gp
Keywords: Adiabatic approximation, quasi-Bell states, reduced density matrices, entropy, $Q$-function.

# I Introduction

One of the simplest and most important models in quantum physics describes a two-level system (qubit) interacting with a simple harmonic oscillator. The system is well-understood [1] in the context of the rotating wave approximation that holds for a small detuning *i.e.* near-equality of the oscillator and the qubit frequency, and a simultaneously small qubit-oscillator coupling compared to the oscillator frequency. A property of this approximation is the conservation of the total number of excitations. In this regime the system is exactly solvable [1] allowing closed-form evaluation of dynamical quantities. Recently, however, a variety of experimental situations pertaining to stronger coupling domain, where the rotating wave approximation is no longer valid, have been investigated. Superconducting qubits interacting with nanomechanical resonators [2-4] is one such promising area where experimentally the qubit-oscillator coupling constant may be driven to values comparable to the qubit and oscillator energy scales. Incorporating terms in the qubit-oscillator interaction Hamiltonian the authors of Ref. [5, 6] advanced an adiabatic approximation scheme that utilizes the separation of the time scales of the qubit and the oscillator. The dynamical state of the fast-moving component, say the oscillator, is assumed to instantaneously adjust to the slow-changing state of the qubit. This allows for decoupling of the full Hamiltonian into sectors related to each time scale, and approximate eigenstates of the system may be derived. The problem has also been studied by using a generalization of the rotating wave approximation [7] where the energy eigenstates have been investigated. For a single qubit analytical expression of the time-dependent behaviour of the two level system was obtained [5] in the small coupling limit.Studying a system of two qubits simultaneously coupled to a single oscillator mode the authors of Ref. [8] analytically evaluated the collapse and revival dynamics in the moderately strong coupling regime. Their procedure involved using Poisson sum formula [9] that leads to expressions of the reduced density matrix of the qubits as an infinite sum.

It has been noted in [6, 10] that whereas in the case of cavity electrodynamics the usual experimental set up is described by an unbiased qubit operated through its degeneracy point, the static bias of a superconducting qubit may be easily varied, say, by operating a magnetic flux on a Josephson junction. The energy spectrum of the system in the strong coupling regime in the presence of a bias has been investigated [10] by employing the Van Vleck perturbation theory [11]. Using a semiclassical approach the ground state in the presence of a bias has been investigated in [6]. Recently the effective Hamiltonian of two superconducting qubits coupled to a oscillator in the presence of a static bias has been explored [12].

In the current work we investigate the moderately strong qubit-oscillator coupling constant regime in the presence of a dominant oscillator frequency and a static bias of the qubit. In particular, starting with a quasi-Bell bipartite initial state we use the adiabatic approximation to find the time-evolution of the reduced density matrices of the qubit and the oscillator, respectively. Retaining up to the quadratic terms in the Laguerre polynomials in the phase factors the density matrix elements may be analytically expressed in closed forms via the Jacobi theta functions. Using this structure we explicitly obtain the entropy of the reduced qubit density matrix, which measures the time-variation of the entanglement of the bipartite state. To evaluate the analytical expressions we treat the bias perturbatively, and retain terms of second order in smallness. The summation procedure may, however, be extended to arbitrary order. Moreover, in the course of our analytical derivation we neglect the bias terms in the exponents of the phase factors as that will be of importance in the asymptotically long time limit. Here we restrict our attention to a finite number of revivals. The reduced density matrix of the oscillator yields the time evolution of the Husimi $Q$-function, and this phase space distribution is subsequently used to analytically evaluate averages of the antinormally ordered operators.

# II The reduced density matrices

The boson-qubit Hamiltonian considered in our work reads, in natural units ($\hbar = 1$), as follows:

$$H = -\frac{\Delta}{2}\sigma_x - \frac{\epsilon}{2}\sigma_z + \omega a^\dagger\, a + \lambda\sigma_z\,(a^\dagger + a), \tag{2.1}$$

where $a^\dagger$ and $a$ are the raising and lowering operators of the harmonic oscillator with frequency $\omega$, the Pauli spin operators $\sigma_x$ and $\sigma_z$ describe the qubit with energy splitting $\Delta$ as well as static bias $\epsilon$, and the qubit-oscillator coupling strength is denoted by $\lambda$. The standard Fock states $\{|n\rangle,\, n = 0, 1, \ldots;\; a\,|n\rangle = \sqrt{n}\,|n-1\rangle, a^\dagger\,|n\rangle = \sqrt{n+1}\,|n+1\rangle\}$ provide the basis for the oscillator, and the following eigenstates $\sigma_z| \pm 1\rangle = \pm| \pm 1\rangle$ span the space of the qubit. The Hamiltonian (2.1) is not known to be exactly solvable. It departs from the rotating wave approximation that holds good in the near-resonance and weak coupling regime. In the present work we follow, as mentioned earlier, the adiabatic approximation [5] that relies on the separation of the time scales characterizing the high oscillator frequency and the (renormalized) low qubit frequency. The fast-moving oscillator then adiabatically adjusts to the slow changes of the state of the qubit.

To proceed one notes that, in the said time scale, the oscillator follows the instantaneous state of the qubit observable $\sigma_z$, and for the oscillator degree of freedom the spin variable $\sigma_z$ may be assumed to be in its eigenstate with the eigenvalue given by $\pm 1$. The effective Hamiltonian of the harmonic oscillator now reads

$$H_{\mathcal{O}} = \omega a^\dagger a \pm \lambda(a^\dagger + a). \tag{2.2}$$

The number states shifted symmetrically by the displacement operator diagonalize the above Hamiltonian:

$$|n_\pm\rangle = \mathrm{D}^\dagger\,(\pm\lambda/\omega)\,|n\rangle, \qquad \mathrm{D}(\alpha) = \exp(\alpha a^\dagger - \alpha^* a), \tag{2.3}$$

where the degenerate eigenenergies are as follows:

$$E_n^\pm \equiv \mathcal{E}_n = n\omega - \frac{\lambda^2}{\omega}. \tag{2.4}$$

The number states displaced in the same direction maintain orthonormality, whereas the overlap between the Fock states displaced in opposite directions is given by

$$\langle m_-|n_+\rangle = \begin{cases} (-1)^{m-n}\ \exp\left(-\frac{x}{2}\right) x^{\frac{m-n}{2}} \sqrt{n!/m!}\ L_n^{m-n}(x)\,, & m \geqslant n \\ \exp\left(-\frac{x}{2}\right) x^{\frac{n-m}{2}} \sqrt{m!/n!}\ L_m^{n-m}(x)\,, & m < n, \end{cases} \tag{2.5}$$

where the associated Laguerre polynomial reads $L_n^j(x) = \sum_{k=0}^{n} (-1)^k \binom{n+j}{n-k} \frac{x^k}{k!}$, and the parameter $x = (2\lambda/\omega)^2$ acts as the perturbative expansion coefficient.

After diagonalizing the high frequency oscillator component of the Hamiltonian one now turns attention to the low frequency qubit part. Tensoring the qubit states with the displaced oscillator basis $\{|\pm 1, n_{\pm}\rangle \,|\, \forall n = 0, 1, \ldots\}$ the Hamiltonian may now be readily constructed. In the parameter regime $\lambda < \omega$ one may neglect the matrix elements that mixes oscillator states with different eigenvalues of its number operator. This reduces the Hamiltonian to a block-diagonal form where each block mixes displaced oscillator states with identical number of photonic excitations. The Hamiltonian for the $n$-th block assumes the form

$$H = \begin{pmatrix} n\omega - \dfrac{\lambda^2}{\omega} - \tilde{\epsilon} & \delta_n \\ \delta_n & n\omega - \dfrac{\lambda^2}{\omega} + \tilde{\epsilon} \end{pmatrix}, \quad \delta_n = -\frac{\widetilde{\Delta}}{2} L_n(x), \; \widetilde{\Delta} = \Delta \exp\left(-\frac{x}{2}\right), \; \tilde{\epsilon} = \frac{\epsilon}{2} \tag{2.6}$$

that may be diagonalized in the basis of the eigenstates listed below

$$|\mathcal{E}_n^{\pm}\rangle = \sqrt{\frac{\chi_n \mp \tilde{\epsilon}}{2\chi_n}} |1, n_+\rangle \pm \frac{\delta_n}{|\delta_n|} \sqrt{\frac{\chi_n \pm \tilde{\epsilon}}{2\chi_n}} |-1, n_-\rangle, \tag{2.7}$$

where the corresponding eigenvalues read

$$\mathcal{E}_n^{\pm} = n\omega - \lambda^2/\omega \pm \chi_n, \qquad \chi_n = \sqrt{\delta_n^2 + \tilde{\epsilon}^2}. \tag{2.8}$$

In order to understand the qubit-oscillator entanglement we study the dynamical evolution of the quasi-Bell initial states of the coupled system in the presence of a bias ($\epsilon \neq 0$):

$$|\psi(0)\rangle^{(\pm)} = \frac{1}{\sqrt{2}} \left(|1, \alpha\rangle \pm |-1, -\alpha\rangle\right), \qquad |\alpha\rangle = \mathrm{D}(\alpha)|0\rangle, \tag{2.9}$$

where the coherent state $\{|\alpha\rangle \; \forall \alpha = \Re(\alpha) + i\Im(\alpha) \in \mathbb{C}\}$ for the oscillator degree of freedom may be realized by the action of the displacement operator on the vacuum. The nonclassical entangled states (2.9) have been discussed earlier [13] in the context of generating bipartite and multipartite entanglements in cavity quantum electrodynamics. It has also been observed [6] that in the large coupling regime $\big(\lambda \gtrsim \omega\big)$ a state of the generic type (2.9) becomes the approximate ground state of the combined system. In the present work we, as an example of a typical nonclassical state, analytically study the properties such as time evolution of its entanglement measure and the phase space distributions of the oscillator for moderately strong qubit-oscillator coupling. The precise limits of the coupling constant and the bias parameter will be made explicit later. The time evolution of the initial state (2.9) is given by

$$\begin{aligned} |\psi(t)\rangle^{(\pm)} \;=\; & \frac{1}{\sqrt{2}} \exp\left(-\frac{1}{2}\varrho_{\hat{\alpha}}^2 - i\left(\Im(\alpha)\frac{\lambda}{\omega} + \mathcal{E}_n t\right)\right) \times \\ & \times \sum_{n=0}^{\infty} \frac{1}{\sqrt{n!}} \widehat{\alpha}^n \left[\Xi_n^{\pm} \exp(-i\chi_n t)|\mathcal{E}_n^{+}\rangle \mp (-1)^n \frac{\delta_n}{|\delta_n|} \; \Xi_n^{\mp} \exp(i\chi_n t)|\mathcal{E}_n^{-}\rangle\right], \end{aligned} \tag{2.10}$$

where $\widehat{\alpha} = \alpha + \lambda/\omega$, $\varrho_{\widehat{\alpha}} = |\widehat{\alpha}|$, and $\Xi_n^{\pm} = \sqrt{\dfrac{\chi_n - \tilde{\epsilon}}{2\chi_n}} \pm (-1)^n \dfrac{\delta_n}{|\delta_n|}\sqrt{\dfrac{\chi_n + \tilde{\epsilon}}{2\chi_n}}$. On the rhs of (2.10) and hereafter the upper and lower signs refer to respective quasi-Bell states given in (2.9). Employing the state (2.10) the evolution of the density matrix of the composite bipartite system may be readily constructed:

$$\rho^{(\pm)}(t) = |\psi(t)\rangle^{(\pm)}\langle\psi(t)|. \tag{2.11}$$

The marginalized density matrices for the individual degrees of freedom may be obtained by partial tracing of the composite density matrix (2.11). For instance the tracing of the oscillator degree of freedom yields the reduced density matrix of the qubit. Its explicit structure obtained via (2.11, 2.10) assumes the form

$$\rho_{\mathcal{Q}}^{(\pm)}(t) \equiv \mathrm{Tr}_{\mathcal{O}}\,\rho^{(\pm)}(t) = \begin{pmatrix} \dfrac{1}{2} \mp \zeta & \pm\xi \\ \pm\xi^{*} & \dfrac{1}{2} \pm \zeta \end{pmatrix}, \tag{2.12}$$

where the components read

$$\begin{aligned} \zeta &= \tilde{\epsilon}\, \exp\left(-\varrho_{\widehat{\alpha}}^{2}\right) \sum_{n=0}^{\infty} (-1)^n \frac{\varrho_{\widehat{\alpha}}^{2n}}{n!}\, \delta_n\, \frac{\sin^2 \chi_n t}{\chi_n^2}, & (2.13) \\ \xi &= \frac{1}{2}\, \exp(-\varrho_{\widehat{\alpha}}^{2}) \sum_{n=0}^{\infty} (-1)^n \frac{\varrho_{\widehat{\alpha}}^{2n}}{n!} \left(1 - 2\tilde{\epsilon}^2\, \frac{\sin^2 \chi_n t}{\chi_n^2} + i\,\tilde{\epsilon}\, \frac{\sin 2\chi_n t}{\chi_n}\right) \langle n_-|n_+\rangle. & (2.14) \end{aligned}$$

An interesting feature evident in the reduced density matrix (2.12) that it is *time-independent* in the limit $\epsilon = 0$. Physically one may conclude that when the fast-changing oscillator degree of freedom is traced out the slow-moving qubit degree of freedom is close to its maximally mixed form that becomes time-independent in the bias-free limit.

Similarly the trace over the qubit basis generates the reduced density matrix of the oscillator:

$$\rho_{\mathcal{O}}^{(\pm)}(t) \equiv \mathrm{Tr}_{\mathcal{Q}}\,|\psi(t)\rangle^{(\pm)}\langle\psi(t)| \tag{2.15}$$

that may be expressed, via (2.10), in terms of the displaced number states as follows:

$$\begin{aligned} \rho_{\mathcal{O}}^{(\pm)}(t) &= \frac{1}{2}\exp(-\varrho_{\widehat{\alpha}}^{2}) \sum_{n,m=0}^{\infty} \frac{(\widehat{\alpha})^n\,(\widehat{\alpha}^{*})^m}{\sqrt{n!m!}} \Big(\mathfrak{C}_n^{(\pm)}(t)\mathfrak{C}_m^{(\pm)^{*}}(t)|n_+\rangle\langle m_+| \\ &\quad +(-1)^{n+m}\mathfrak{C}_n^{(\mp)^{*}}(t)\mathfrak{C}_m^{(\mp)}(t)|n_-\rangle\langle m_-|\Big) \exp\big(-i(n-m)\omega t\big), \end{aligned} \tag{2.16}$$

where the coefficients read

$$\mathfrak{C}_n^{(\pm)}(t) = \mathfrak{A}_n^{(\mp)} \exp(i\chi_n t) + \mathfrak{B}_n^{(\pm)} \exp(-i\chi_n t), \tag{2.17}$$

$$\mathfrak{A}_n^{(\pm)} = \frac{\chi_n + \tilde{\epsilon} \pm (-1)^n \delta_n}{2\chi_n}, \quad \mathfrak{B}_n^{(\pm)} = \frac{\chi_n - \tilde{\epsilon} \pm (-1)^n \delta_n}{2\chi_n}. \tag{2.18}$$

We note that apart from studying in the context of adiabatic approximation no other simplification has been made in the derivation leading to the reduced density matrix elements (2.13, 2.14, 2.16).

# III The qubit density matrix

We now focus our attention towards analytical evaluation of the density matrix in the parameter regime where the oscillator frequency sets the dominant scale and the following hierarchy is maintained: $\epsilon < \Delta < \omega$. In the weak coupling limit $\lambda \ll \omega$ the series in (2.13, 2.14) may be summed by retaining *à la* [5] only linear terms in the Laguerre polynomials: $L_n(x) = 1 - nx + O(x^2)$. For simplicity we retain terms $O(\epsilon^2)$ in the bias parameter. Since we are interested in the first few revivals of the system we also systematically neglect $\epsilon$ dependent terms in phases as they will be important in very large time limit. The higher order terms in the bias parameter $\epsilon$ may, however, be taken into account in the present perturbative scheme. In this parametric domain the elementary trigonometric identities now allow us to compute the diagonal and the off-diagonal elements of the density matrix given in (2.12-2.14) in the following form:

$$
\begin{aligned}
\zeta &= -\frac{\epsilon}{2\widetilde{\Delta}}\,(1-f)\,\exp\left(-2\varrho^2_{\hat{\alpha}}\right) + \frac{\epsilon}{2\widetilde{\Delta}}\exp\left(-2\varrho^2_{\hat{\alpha}}\cos^2\frac{x\tau}{2}\right)\left(\cos\widehat{\Omega}_t - f\,\cos\widehat{\Omega}^{(x)}_t\right), \quad (3.1)\\
\xi &= \frac{1+f}{2}\exp\left(-2\varrho^2_{\hat{\alpha}} - \frac{x}{2}\right) - \frac{\epsilon^2}{2\Delta\widetilde{\Delta}}\Big[(1-f)\exp\left(-2\varrho^2_{\hat{\alpha}}\right) - \exp\Big(-2\varrho^2_{\hat{\alpha}}\cos^2\frac{x\tau}{2}\Big)\times\\
&\quad\times\left(\cos\widehat{\Omega}_t - f\cos\widehat{\Omega}^{(x)}_t\right)\Big] + i\frac{\epsilon}{2\Delta}\exp\left(-2\varrho^2_{\hat{\alpha}}\cos^2\frac{x\tau}{2}\right)\sin\widehat{\Omega}_t, \quad (3.2)
\end{aligned}
$$

where the scaled time variable as well as the phases are given, respectively, by $\tau = \widetilde{\Delta}\,t, \widehat{\Omega}_t = \varrho^2_{\hat{\alpha}}\sin(x\tau) + \tau$, $\widehat{\Omega}^{(x)}_t = \varrho^2_{\hat{\alpha}}\sin(x\tau) + (1-x)\,\tau$, and the parameter $f = x\varrho^2_{\hat{\alpha}}$ determines the amplitude of the mode with the renormalized frequency. This approximation that retains up to linear terms in Laguerre polynomials hold in the weak coupling range. For higher qubit-oscillator coupling the linear approximation method fails for the higher revival times.

In an attempt for analytically evaluating the density matrix elements in the context of Tavis-Cummings model for higher values of the qubit-oscillator coupling constant the authors of Ref. [8] considered a method based on Poisson sum formula. Here one retains contributions of the order $O(x^2) \sim O((\lambda/\omega)^4)$ that amounts to considering up to the quadratic terms in the expansion of the Laguerre polynomials: $L_n(x) = 1 - nx + \frac{1}{2!}\binom{n}{2}x^2 + O(x^3)$. Following [8] we now provide a Poisson sum for the elements of the density matrix (2.12). To proceed, for a large value of $\varrho^2_{\hat{\alpha}}$, we approximate the Poisson distribution in (2.13, 2.14) with the corresponding Gaussian limit:

$$
P(n) \equiv \exp\left(-\varrho^2_{\hat{\alpha}}\right)\frac{\varrho^{2n}_{\hat{\alpha}}}{n!} \to \frac{1}{\sqrt{2\pi\varrho^2_{\hat{\alpha}}}}\exp\left(-\frac{\left(n-\varrho^2_{\hat{\alpha}}\right)^2}{2\varrho^2_{\hat{\alpha}}}\right). \quad (3.3)
$$

The Poisson summation formula employs Fourier series and, in the present context, it may be stated [9] as follows: An arbitrary function $\mathfrak{g}(\mathsf{x})$ integrable in the domain $(-\infty, \infty)$ satisfies the sum rule

$$
\sum_{n=-\infty}^{\infty}\frac{\mathfrak{g}(n^+) + \mathfrak{g}(n^-)}{2} = \sum_{\ell=-\infty}^{\infty}\int_{-\infty}^{\infty}\mathfrak{g}(\mathsf{t})\exp(-2\pi i\ell\mathsf{t})\,d\mathsf{t}. \quad (3.4)
$$

For the case $\varrho^2_{\hat{\alpha}} \gg 1$ the lower limit of integration corresponding to (3.3) may be extended to $(-\infty)$, and the integrals in the Fourier transforms may be evaluated by the standard

Gaussian integration technique. With the above structure in mind we obtain the Poisson sum for the diagonal term (2.13) in the density matrix:

$$\begin{aligned}\zeta &= -\frac{\epsilon}{2\widetilde{\Delta}}\exp\left(-2\varrho_{\widehat{\alpha}}^{2}\right)\left(1-f(1+x)+\frac{3}{4}f^{2}\right)\\ &\quad+\frac{\epsilon}{2\widetilde{\Delta}}\sqrt{\cos\theta}\sum_{\ell=0}^{\infty}\Re\big[\mathcal{I}_{\ell}^{(1)}-f(1+x)\,\mathcal{I}_{\ell}^{(2)}+\frac{3}{4}f^{2}\,\mathcal{I}_{\ell}^{(3)}\big].\end{aligned}\tag{3.5}$$

In (3.5) the terms comprising the Fourier components read

$$\begin{aligned}\mathcal{I}_{\ell}^{(\jmath)}\Big|_{\jmath\in\{1,2,3\}} &= \exp\left(-\varsigma^{(\jmath)}\left(\tau-\upsilon_{\ell}^{(\jmath)}\right)^{2}\right)\quad\times\\ &\quad\times\exp\left[i\left\{\varsigma\left(\tan\theta\left(\left(\varphi_{\ell}^{(\jmath)}(\tau)\right)^{2}-1\right)+2\varphi_{\ell}^{(\jmath)}(\tau)\right)-\tau^{(\jmath)}-\frac{\theta}{2}\right\}\right],\end{aligned}\tag{3.6}$$

where the parameters are given by $\tan\theta=\dfrac{xf\tau}{2}, \varsigma=\dfrac{1}{2}\,\varrho_{\widehat{\alpha}}^{2}\,\cos^{2}\theta$. The strength of the Gaussian packets in the rhs of (3.5) on the respective powers of the parameter $f$. The values of the scaled time variable $\tau$ at which the peak of the contributions (3.6) occur, and the decay of the corresponding exponentials are given, respectively, by

$$\upsilon_{\ell}^{(\jmath)}=\frac{2\pi(\ell+1/2)}{x\left(1+\dfrac{(3-2\jmath)x}{4}-\dfrac{f}{2}\right)},\quad \varsigma^{(\jmath)}=\varsigma x^{2}\left(1+\frac{(3-2\jmath)x}{4}-\frac{f}{2}\right)^{2}.\tag{3.7}$$

The phase constructs in (3.6) read

$$\varphi_{\ell}^{(\jmath)}(\tau)=x\left(1+\frac{(3\jmath-2)x}{4}\right)\tau-2\pi(\ell+1/2)\tag{3.8}$$

and the scaled time-dependent parameters are given by

$$\tau^{(1)}=\tau,\ \tau^{(2)}=(1-x)\tau,\ \tau^{(3)}=\left(1-2x+\frac{x^{2}}{2}\right)\tau.\tag{3.9}$$

We mention that in evaluating the first term in the rhs of (3.5) the Gaussian limit (3.3) has not been exploited as the corresponding term in (2.13) is free from a complex Laguerre polynomial dependent phase and is amenable to elementary summation technique. Similarly, in summing other series expressions we will continue to use elementary procedures wherever applicable.

Continuing the above Poisson sum process the off-diagonal element (2.14) for the reduced density matrix of the qubit may be evaluated as

$$\begin{aligned}\xi &= \frac{1}{2}\exp\left(-2\varrho_{\widehat{\alpha}}^{2}-x/2\right)\left(1+f+\frac{f^{2}}{4}\right)-\frac{\epsilon^{2}}{2\Delta\widetilde{\Delta}}\Big[\exp\left(-2\varrho_{\widehat{\alpha}}^{2}\right)\left(1-f(1+x)+\frac{3}{4}f^{2}\right)\\ &\quad-\sqrt{\cos\theta}\sum_{\ell=0}^{\infty}\Re\big[\mathcal{I}_{\ell}^{(1)}-f(1+x)\mathcal{I}_{\ell}^{(2)}+\frac{3}{4}f^{2}\mathcal{I}_{\ell}^{(3)}\big]\Big]-\frac{i\epsilon}{2\Delta}\sqrt{\cos\theta}\sum_{\ell=0}^{\infty}\Im\big[\mathcal{I}_{\ell}^{(1)}\big].\end{aligned}\tag{3.10}$$

The contribution of $\mathcal{I}_{\ell}^{(\jmath)}$ in the reduced density matrix elements (3.5, 3.10), as evident from (3.6), is maximum when its real part achieves maximum value at $\tau = \upsilon_{\ell}^{(\jmath)}$ that corresponds to successive revivals of the two-level system. The index $\ell$, therefore, has the natural significance of enumerating specific revivals of the system. A sharp peak corresponds to higher value of the fall-off parameter $\varsigma^{(\jmath)}$. From the structure of (3.6) it follows that for a positive $\tau$ only the Fourier components with positive indices $\ell \geqslant 0$ need to be summed over, as it was maintained in obtaining the density matrix elements (3.5, 3.10).

In the said limit of moderately strong coupling regime where we retain contributions $O(x^2)$ in the density matrix elements and consider up to quadratic terms in the Laguerre polynomials an alternate method of summing the series in (2.13, 2.14) may be proposed by employing the Jacobi theta functions. This method has the additional benefit in that it expresses the matrix elements in closed form. We quote [9] the series expansions of the Jacobi theta function that is relevant in our case:

$$\vartheta_3(q,z) = \sum_{n=-\infty}^{\infty} q^{n^2} \exp(2inz), \qquad \vartheta_4(q,z) = \sum_{n=-\infty}^{\infty} (-1)^n q^{n^2} \exp(2inz). \tag{3.11}$$

These interrelated functions $\vartheta_3(q,z), \vartheta_4(q,z)$ depend on two complex arguments and the series sums in (3.11) are convergent for the choice $|q| < 1$. Our starting point for evaluating the infinite sums in the series (2.13, 2.14) via the theta functions is that we again use the Gaussian limit expressed in (3.3). For a large value of $\varrho_{\hat{\alpha}}^2$ the lower limit of the series in (2.13, 2.14) may be extended to $(-\infty)$. Due to the sharp decay of the Gaussian function the error encountered is quite small. This process is similar to the evaluation of Fourier components via the Gaussian integration technique in the previously described Poisson-sum method, where the lower limit of integration is extended to $(-\infty)$. We here mention that a truncated series on one side in the definition (3.11) remains convergent in the domain $|q| < 1$, and may be viewed as a close analogue of the theta function. Since the difference between the alternate cases in our parametric regime is quite small we retain the standard definition (3.11). The resultant closed-form expression for the diagonal term (2.13) now assumes the form:

$$\zeta = -\frac{\epsilon}{2\widetilde{\Delta}}\Big[\exp\Big(-2\varrho_{\hat{\alpha}}^2\Big)\Big(1 - f(1+x) + \frac{3}{4}f^2\Big) - \frac{\exp\Big(-\varrho_{\hat{\alpha}}^2/2\Big)}{\sqrt{2\pi\varrho_{\hat{\alpha}}^2}}\Re\big(\Theta(\tau)\big)\Big], \tag{3.12}$$

where $\Theta(\tau)$ is explicitly given as a sum of theta functions as follows:

$$\begin{aligned} \Theta(\tau) &= \exp\left(-i\tau^{(1)}\right)\vartheta_4(q, z^{(1)}) - f(1+x)\exp\left(-i\tau^{(2)}\right)\vartheta_4(q, z^{(2)}) \\ &\quad + \frac{3}{4}f^2\exp\left(-i\tau^{(3)}\right)\vartheta_4(q, z^{(3)}). \end{aligned} \tag{3.13}$$

The three theta functions appearing in the combination (3.13) posses same value of the complex argument $q$ but differing values of the other complex variable given, respectively, as

$$q(\tau) = \exp\Big(-\frac{1}{2\varrho_{\hat{\alpha}}^2}\Big)\exp\Big(-i\frac{x^2\tau}{4}\Big), \quad z^{(\jmath)}(\tau)\Big|_{\jmath\in\{1,2,3\}} = \frac{1}{2}\Big(x\Big(1 + (3-2\jmath)\frac{x}{4}\Big)\tau - i\Big). \tag{3.14}$$

For the off-diagonal element (2.14) a similar closed-form expression involving theta functions may also be obtained as follows:

$$
\begin{aligned}
\xi &= \frac{1}{2}\left(1+f+\frac{f^2}{4}\right)\exp\left(-2\varrho_{\hat{\alpha}}^2-\frac{x}{2}\right)-\frac{\epsilon^2}{2\Delta\widetilde{\Delta}}\Big[\exp\big(-2\varrho_{\hat{\alpha}}^2\big) \quad \times \\
&\quad \times\left(1-f(1+x)+\frac{3}{4}f^2\right)-\frac{\exp\big(-\varrho_{\hat{\alpha}}^2/2\big)}{\sqrt{2\pi\varrho_{\hat{\alpha}}^2}}\Re\big[\Theta(\tau)\big]\Big] \\
&\quad -i\frac{\epsilon}{2\Delta}\frac{\exp\big(-\varrho_{\hat{\alpha}}^2/2\big)}{\sqrt{2\pi\varrho_{\hat{\alpha}}^2}}\Im\big[\exp\left(-i\tau^{(1)}\right)\vartheta_4(q,z^{(1)})\big]. \qquad (3.15)
\end{aligned}
$$

This completes our procedure of evaluations of the elements of the reduced density matrix of the qubit via the Jacobi theta functions. It involves considering up to the quadratic terms in the expression of Laguerre polynomials appearing in the arguments of the phase factors and also approximating the Poisson term with its Gaussian limit. For moderately large coupling ($\lambda/\omega \sim 0.25$) this summation technique has good agreement with the numerical evaluations based on (2.13, 2.14). Concerning our choice of bias parameter we note that as we have included terms up to $O(\epsilon^2)$, we may maintain the order of the dimensionless ratio $\frac{\epsilon}{\widetilde{\Delta}} \sim 0.1$. These restrictions describe the parametric regime in which our analytical evaluations based on theta functions hold. In Figs. [1-3] we now compare the closed-form expressions of the density matrix elements (3.12, 3.15) with the corresponding sums given in (2.13, 2.14). For a significant number of revivals good agreement is observed.



Figure 1: The diagonal component $\zeta$ of the density matrix for (i) $\lambda = 0.1\omega, \Delta = 0.15\omega, \epsilon = 0.01\omega$ and $\alpha = 2$ (left) and (ii) $\lambda = 0.18\omega, \Delta = 0.3\omega, \epsilon = 0.02\omega$ and $\alpha = 1$ (right) using (a) the series (2.13), and (b) the evaluation (3.12) using theta function.



Figure 2: The real part of the off-diagonal component $\xi$ for (i) $\lambda = 0.1\omega, \Delta = 0.15\omega, \epsilon = 0.01\omega$ and $\alpha = 2$ (left) and (ii)$\lambda = 0.18\omega, \Delta = 0.3\omega, \epsilon = 0.02\omega$ and $\alpha = 1$ (right) using (a) the series (2.14), and (b) the evaluation (3.15) using theta function.

The time evolution of the entanglement of the initially chosen quasi-Bell state (2.9) is of interest. For the bipartite system considered here this may be measured by the entropy of the reduced density matrix of the qubit: $S(\rho_Q) = -\mathrm{Tr}(\rho_Q \ln \rho_Q)$. The construction (2.12) of the reduced density matrix $\rho_Q$ produces its pair of eigenvalues given below:

$$\Lambda^{\pm} = \frac{1}{2} \pm \sqrt{\zeta^2 + |\xi|^2}, \tag{3.16}$$

which are valid for both the initial choices (2.9) of quasi-Bell states. The said entropy now reads

$$S = -\Lambda^{+} \log \Lambda^{+} - \Lambda^{-} \log \Lambda^{-}. \tag{3.17}$$

The structure of the diagonal and the off-diagonal terms given in (2.13, 2.14) may now be employed to extract the eigenvalues of the density matrix (2.12) up to the order $O(\epsilon^2)$:

$$\Lambda^{\pm} = \frac{1}{2}\Big(1 \pm \exp(-\varrho_{\hat{\alpha}}^2 - x/2)\sqrt{\mathfrak{S}}\Big), \tag{3.18}$$

where the sum is given by

$$\begin{aligned} \mathfrak{S} \;&=\; \sum_{n,m=0}^{\infty} (-1)^{(n+m)} \frac{\varrho_{\hat{\alpha}}^{2(n+m)}}{n!m!} L_n(x) L_m(x) \quad \times \qquad (3.19) \\ &\times \;\; \Big(1 - \epsilon^2 \frac{\sin^2(\chi_n t)}{\chi_n^2} + \epsilon^2 \frac{\sin(2\chi_n t)}{2\chi_n} \frac{\sin(2\chi_m t)}{2\chi_m} + \epsilon^2 \Big(\frac{\Delta}{2}\Big)^2 \frac{\sin^2(\chi_n t)}{\chi_n^2} \frac{\sin^2(\chi_m t)}{\chi_m^2}\Big) . \end{aligned}$$



Figure 3: The imaginary part of the off-diagonal component $\xi$ for (i) $\lambda = 0.1\omega, \Delta = 0.15\omega, \epsilon = 0.01\omega$ and $\alpha = 2$ (left) and (ii)$\lambda = 0.18\omega, \Delta = 0.3\omega, \epsilon = 0.02\omega$ and $\alpha = 1$ (right) using (a) the series (2.14), and (b) the evaluation (3.15) using theta function.

It is interesting to consider the entropy in the bias-free $\epsilon = 0$ limit. Towards this end we use the identity

$$\sum_{n=0}^{\infty} \frac{(-1)^n}{n!} L_n(\mathsf{x})\mathsf{t}^n = \exp(-\mathsf{x}) I_0(2\sqrt{\mathsf{xt}}), \tag{3.20}$$

where the Bessel function is denoted as

$$I_0(z) = \sum_{k=0}^{\infty} \frac{(z/2)^{2k}}{(k!)^2}. \tag{3.21}$$

The equations ranging (3.17-3.21) may now be used to obtain an exact closed-form expression for the entropy in the bias-free limit:

$$S\,|_{\epsilon=0} = \ln 2 - \frac{1}{2} \exp\big(-4\varrho_{\hat{\alpha}}^2 - x\big)\Big(I_0\big(2\sqrt{f}\big)\Big)^2. \tag{3.22}$$

The entropy (3.22) in the $\epsilon = 0$ limit is independent of time and attains its maximal value for a large $\varrho_{\hat{\alpha}}$.

It is evident from (3.19) that the inclusion of a bias term modifies the entropy (3.22) only at the order $O(\epsilon^2)$. We first obtain the entropy in the weak coupling limit where the linear approximation of the Laguerre functions as advocated in [5] is assumed to be valid. Our previous constructions given in (3.1, 3.2, 3.17, 3.19) now produce the entropy in the said limit:

$$\begin{aligned} S &= \ln 2-\frac{1}{2}\,\exp(-4\varrho_{\widehat{\alpha}}^2-x)\,\Big(1+2f-\frac{2\epsilon^2}{\widetilde{\Delta}^2}\,\Big(1-\Big((1+f)\cos\widehat{\Omega}_t-f\cos\widehat{\Omega}_t^{(x)}\Big)\;\times \\ &\quad\times\exp\Big(2\varrho_{\widehat{\alpha}}^2\sin^2\frac{x\tau}{2}\Big)-\frac{1}{2}\exp\Big(4\varrho_{\widehat{\alpha}}^2\sin^2\frac{x\tau}{2}\Big)\sin^2\widehat{\Omega}_t-\frac{1}{2}\exp(x)(1-2f) \\ &\quad-\frac{1}{2}\exp\Big(4\varrho_{\widehat{\alpha}}^2\sin^2\frac{x\tau}{2}+x\Big)\Big(\cos^2\widehat{\Omega}_t-2f\cos\Omega_t\cos\Omega_t^{(x)}\Big)\Big)\Big)\,. \end{aligned} \tag{3.23}$$

It is easily evident that the expression of entropy (3.23) in the weak coupling limit agrees in the bias-free $\epsilon = 0$ case with the evaluation (3.22) when the rhs of the latter is expanded retaining up to terms linear in the parameter $f$.

For the moderately strong coupling described by the parametric regime specified in the context of the equation (3.15) we evaluate the entropy (3.17) by retaining terms up to the order $O(x^2)$ in the Laguerre polynomials. The summation procedure as evidenced in (3.12, 3.15) now yields the entropy via linear combinations of Jacobi theta functions. Up to the order in the bias parameter $O(\epsilon^2)$ the entropy reads

$$\begin{aligned} S &= \ln 2-\frac{1}{2}\,\exp\big(-4\varrho_{\widehat{\alpha}}^2-x\big)\Big[1+2f+\frac{3}{2}f^2-\frac{2\epsilon^2}{\widetilde{\Delta}^2}\Big[1-fx-f^2x-\Big(1+f+\frac{f^2}{4}\Big)\;\times \\ &\quad\times\;\frac{\exp\big(3\varrho_{\widehat{\alpha}}^2/2\big)}{\sqrt{2\pi\varrho_{\widehat{\alpha}}^2}}\Re\big(\Theta(\tau)\big)-\frac{\exp(3\varrho_{\widehat{\alpha}}^2)}{4\pi\varrho_{\widehat{\alpha}}^2}\Big(\Im\big(\exp\big(-i\tau^{(1)}\big)\,\vartheta_4(q,z^{(1)})\big)\Big)^2 \\ &\quad+\frac{\exp\big(3\varrho_{\widehat{\alpha}}^2/2+x\big)}{\sqrt{2\pi\varrho_{\widehat{\alpha}}^2}}\Big(1-f(1+x)+\frac{3}{4}f^2\Big)\,\Re\big(\Theta(\tau)\big) \\ &\quad-\frac{\exp\big(3\varrho_{\widehat{\alpha}}^2+x\big)}{4\pi\varrho_{\widehat{\alpha}}^2}\Big(\Re\big(\Theta(\tau)\big)\Big)^2-\frac{\exp(x)}{2}\Big(1-2f(1+x)+f^2\Big(\frac{5}{2}+2x\Big)\Big)\Big]\Big]. \end{aligned} \tag{3.24}$$

We present below in Fig. [4] the plots of the time evolution of the entropy of the system based on our closed-form expression (3.24) and compare it with sum given in (3.17, 3.19). For first few revivals good agreement is obtained. As physically expected the entropy dips from its maximal value during successive revivals of the system. The extent of the dip increases with the increasing bias parameter. For the bias-free case ($\epsilon = 0$) the entropy remains constant.

# IV The $Q$-function of the oscillator density matrix

The Husimi $Q$-function [14] is one of the distribution functions that facilitates characterization of the quantum state of the oscillator degree of freedom. For our choice of quasi-Bell initial states (2.9) it is defined as the expectation value of the reduced density matrix of the oscillator (2.16) in an arbitrary coherent state:

$$Q^{(\pm)}(\beta) = \frac{1}{\pi}\langle\beta|\rho_{\mathcal{O}}^{(\pm)}(t)|\beta\rangle. \tag{4.1}$$



Figure 4: Evaluation of the entropy for (i) $\lambda = 0.1\omega, \Delta = 0.15\omega, \epsilon = 0.01\omega$ and $\alpha = 2$ (left) and (ii) $\lambda = 0.18\omega, \Delta = 0.3\omega, \epsilon = 0.02\omega$ and $\alpha = 1$ (right) using (a) the series (3.17), and (b) the evaluation (3.24) using theta function.

Substituting the said reduced density matrix evaluated in (2.16) the $Q$-function of the state under consideration may be written as,

$$\begin{aligned} Q^{(\pm)}(\beta) &= \frac{1}{2\pi}\exp(-\varrho^2_{\widehat{\alpha}}) \sum_{n,m=0}^{\infty} \Big[ \exp(-\varrho^2_{\widehat{\beta}}) \frac{(\widehat{\alpha}\widehat{\beta}^*)^n\,(\widehat{\alpha}^*\widehat{\beta})^m}{n!m!} \Big(\mathfrak{c}^{(\pm)}_n(t)\mathfrak{c}^{(\pm)*}_m(t) \\ &\quad + (-1)^{n+m}\exp(-\varrho^2_{\breve{\beta}}) \frac{(\widehat{\alpha}\breve{\beta}^*)^n\,(\widehat{\alpha}^*\breve{\beta})^m}{n!m!} \mathfrak{c}^{(\mp)*}_n(t)\mathfrak{c}^{(\mp)}_m(t)\Big)\Big] \quad \times \\ &\quad \times \exp(-i(n-m)\omega t), \end{aligned} \tag{4.2}$$

where we abbreviate: $\widehat{\beta} = \beta + \lambda/\omega, \varrho_{\widehat{\beta}} = |\widehat{\beta}|$ and $\breve{\beta} = \beta - \lambda/\omega, \varrho_{\breve{\beta}} = |\breve{\beta}|$. To evaluate the sum in (4.2) we adopt a procedure discussed in the context of (3.1, 3.2) that, following [5], entails retaining up to linear terms in the expansion of Laguerre polynomials. As the two sums in (4.2) factorizes they may be easily obtained in the said limit. The positive definite $Q$-function now assumes the form

$$Q^{(\pm)}(\beta) = \frac{1}{2\pi}\exp(-\varrho^2_{\widehat{\alpha}})\Big(\exp(-\varrho^2_{\widehat{\beta}})|\mathfrak{X}^{(\pm)}|^2 + \exp(-\varrho^2_{\breve{\beta}})|\mathfrak{Y}^{(\pm)}|^2\Big), \tag{4.3}$$

where the complex density functions read

$$\begin{aligned} \mathfrak{X}^{(\pm)} &= \exp(\Phi^{\|}_t)\cos\left(\Phi^{\perp}_t - \frac{\tau}{2}\right) \pm i\exp(-\Phi^{\|}_t)\sin\left(\Phi^{\perp}_t + \frac{\tau}{2}\right) \\ &\quad -i\frac{\epsilon}{\widetilde{\Delta}}\exp(\Phi^{\|}_t)\left[\sin\left(\Phi^{\perp}_t - \frac{\tau}{2}\right) + x\,\Phi_t\sin\left(\Phi^{\perp}_t - \frac{(1-x)\tau}{2}\right)\right] \\ &\quad \mp i\frac{\epsilon^2}{2\widetilde{\Delta}^2}\exp(-\Phi^{\|}_t)\left[\sin\left(\Phi^{\perp}_t + \frac{\tau}{2}\right) - 2x\Phi_t\sin\left(\Phi^{\perp}_t + \frac{(1-x)\tau}{2}\right)\right], \end{aligned} \tag{4.4}$$



Figure 5: The $Q^{(+)}(\beta)$ function using (i) the series (4.2), and (ii) the analytical result (4.3), in left and right columns respectively, for $\lambda = 0.08\omega, \Delta = 0.15\omega, \epsilon = 0.01\omega$ and $\alpha = 2$ at various time $t$ (a)0 (b)300 (c)500 (d)900.

and

$$\begin{aligned}\mathfrak{Y}^{(\pm)} \;=\;& \exp(-\Psi_t^{\|})\cos\left(\Psi_t^{\perp} + \frac{\tau}{2}\right) \mp i\exp(\Psi_t^{\|})\sin\left(\Psi_t^{\perp} - \frac{\tau}{2}\right) \\ & -i\frac{\epsilon}{\widetilde{\Delta}}\exp(-\Psi_t^{\|})\Big(\sin\left(\Psi_t^{\perp} + \frac{\tau}{2}\right) - x\,\Psi_t \sin\left(\Psi_t^{\perp} + \frac{(1-x)\tau}{2}\right)\Big) \\ & \pm i\frac{\epsilon^2}{2\widetilde{\Delta}^2}\exp(\Psi_t^{\|})\Big(\sin\left(\Psi_t^{\perp} - \frac{\tau}{2}\right) + 2x\,\Psi_t \sin\left(\Psi_t^{\perp} - \frac{(1-x)\tau}{2}\right)\Big).\end{aligned} \tag{4.5}$$

In (4.4, 4.5) we have used the notations $\Phi_t = \widehat{\alpha}\widehat{\beta}^*\exp(-i\omega t)$, $\Psi_t = \widehat{\alpha}\check{\beta}^*\exp(-i\omega t)$, $\Phi_t^{\|} = \Phi_t\cos\frac{x\tau}{2}$, $\Psi_t^{\|} = \Psi_t\cos\frac{x\tau}{2}$, $\Phi_t^{\perp} = \Phi_t\sin\frac{x\tau}{2}$ and $\Psi_t^{\perp} = \Psi_t\sin\frac{x\tau}{2}$. In the parametric region $\lambda/\omega \lesssim 0.1$ where the linear approximation holds we now compare the analytical evaluation (4.3) of the $Q$-function with the corresponding double sum (4.2). The time evolution of the $Q$-function is faithfully reproduced by the closed-form expression (4.3).

One of the utilities of the phase space distribution $Q$ is that it provides a convenient evaluation of the expectation values of the operators expressed in their antinormal ordered form. For instance, the first and second moments of the scaled electric field operator defined as

$$\mathsf{E} = \frac{1}{\sqrt{2}i}(a - a^\dagger) \tag{4.6}$$

may be expressed [14] via the following phase space integrals over the complex plane:

$$\begin{aligned}
\langle \mathsf{E} \rangle^{(\pm)} &\equiv \mathrm{Tr}(\mathsf{E}(a, a^\dagger)\rho_{\mathcal{O}}^{(\pm)}(t)) = \frac{1}{\sqrt{2}i}\int_{-\infty}^{\infty} (\beta - \beta^*)Q^{(\pm)}(\beta)d^2\beta, &(4.7)\\
\langle \mathsf{E}^2 \rangle^{(\pm)} &\equiv \mathrm{Tr}(\mathsf{E}^2(a, a^\dagger)\rho_{\mathcal{O}}^{(\pm)}(t) = -\frac{1}{2}\int_{-\infty}^{\infty} \Big((\beta - \beta^*)^2 + 1\Big)Q^{(\pm)}(\beta)d^2\beta. &(4.8)
\end{aligned}$$

Employing the Fourier sum (4.2) and using the following notation for the coefficient

$$\mathcal{G}_{n;\ell}^{(\pm)}(t) = \widehat{\alpha}^{\ell}\,\Big(\mathfrak{C}_{n+\ell}^{(\pm)}(t)\mathfrak{C}_{n}^{(\pm)*}(t) + (-1)^{\ell}{\mathfrak{C}_{n+\ell}^{(\mp)}}^{*}(t)\mathfrak{C}_{n}^{(\mp)}(t)\Big)\ \exp(-i\ell\omega t) \tag{4.9}$$

the first two moments of the electric field is observed as

$$\begin{aligned}
\langle \mathsf{E} \rangle^{(\pm)} &= \frac{1}{\sqrt{2}}\exp(-\varrho_{\widehat{\alpha}}^2)\sum_{n=0}^{\infty}\frac{\varrho_{\widehat{\alpha}}^{2n}}{n!}\Im\left(\mathcal{G}_{n;1}^{(\pm)}(t)\right), &(4.10)\\
\langle \mathsf{E}^2 \rangle^{(\pm)} &= \varrho_{\widehat{\alpha}}^2 + \frac{1}{2}\Big(1 - \exp(-\varrho_{\widehat{\alpha}}^2)\sum_{n=0}^{\infty}\frac{\varrho_{\widehat{\alpha}}^{2n}}{n!}\,\Re\left(\mathcal{G}_{n;2}^{(\pm)}(t)\right)\Big). &(4.11)
\end{aligned}$$

With the purpose of evaluating the moments (4.10, 4.11) in a closed form we now quote explicit values of the first two Fourier components given in (4.9):

$$\begin{aligned}
\mathcal{G}_{n;1}^{(\pm)}(t) &= \widehat{\alpha}\,\epsilon\Big(\mp(-1)^n\,(\delta_n - \delta_{n+1})\,\frac{\sin\chi_n t}{\chi_n}\,\frac{\sin\chi_{n+1} t}{\chi_{n+1}}\\
&\quad -i\,\left(\frac{\sin\chi_n t}{\chi_n}\,\cos\chi_{n+1}t - \cos\chi_n t\,\frac{\sin\chi_{n+1}t}{\chi_{n+1}}\right)\Big)\,\exp(-i\omega t), &(4.12)\\
\mathcal{G}_{n;2}^{(\pm)}(t) &= 2\widehat{\alpha}^2\Big(\cos\chi_n t\,\cos\chi_{n+2}t + (\tilde{\epsilon}^2 + \delta_n\delta_{n+2})\frac{\sin\chi_n t}{\chi_n}\,\frac{\sin\chi_{n+2}t}{\chi_{n+2}}\\
&\quad \pm i(-1)^n\Big(\delta_n\frac{\sin\chi_n t}{\chi_n}\,\cos\chi_{n+2}t - \delta_{n+2}\,\cos\chi_n t\,\frac{\sin\chi_{n+2}t}{\chi_{n+2}}\Big)\Big)\,\exp(-2i\omega t). &(4.13)
\end{aligned}$$

Utilizing the definitions (3.11) and the construction given in (4.8-4.13) the first two moments of the electric field as linear combinations of of Jacobi $\vartheta$ functions. The first moment is non-vanishing only in the presence of a bias term ($\epsilon \neq 0$) in the qubit Hamiltonian (2.1):

$$\langle \mathsf{E} \rangle^{(\pm)} = -\frac{\epsilon}{\widetilde{\Delta}}\frac{\exp(-\varrho_{\widehat{\alpha}}^2/2)}{\sqrt{4\pi\varrho_{\widehat{\alpha}}^2}}\Big(\Im\,[\,i\widehat{\alpha}\exp(-i\omega t)]\,\Im(\mathsf{T}) \mp\,\Im\,[\widehat{\alpha}\exp(-i\omega t)]\,\Re(\mathsf{H})\Big), \tag{4.14}$$

where the linear combinations of $\vartheta_3$ and $\vartheta_4$ functions read, respectively, as

$$\begin{aligned}
\mathsf{T} &= \left((2 + x + x^2) + \Big(2 + \frac{7x}{2}\Big)f\exp\Big(\frac{-i\tau x^2}{4}\Big) + \frac{3}{2}f^2\exp\Big(\frac{-i\tau x^2}{2}\Big)\right)\times\\
&\quad\times\exp\Big(\frac{i\tau x}{2}\Big)\vartheta_3(\mathfrak{q}, \mathsf{z}) + x(1 + x)\exp\Big(-i\tau\Big(1 - \frac{x}{2}\Big)\Big)\vartheta_3(q, \mathsf{z}^{(1)})\\
&\quad+\frac{3}{2}xf\exp\Big(-i\tau\Big(1 - \frac{3x}{2} + \frac{x^2}{4}\Big)\Big)\,\vartheta_3(q, \mathsf{z}^{(2)}), &(4.15)
\end{aligned}$$

and

$$\begin{aligned}\mathsf{H} \;=\; & \Big(x(1+x) - \frac{3}{2}xf\exp\Big(-\frac{i\tau x^2}{4}\Big)\Big)\exp\Big(\frac{i\tau x}{2}\Big)\vartheta_4(\mathsf{q},\mathsf{z}) \\ & -x(1+x)\exp\Big(-i\tau\Big(1-\frac{x}{2}\Big)\Big)\vartheta_4(q,\mathsf{z}^{(1)}) \\ & +\frac{3}{2}xf\exp\Big(-i\tau\Big(1-\frac{3x}{2}+\frac{x^2}{4}\Big)\Big)\vartheta_4(q,\mathsf{z}^{(2)}). \qquad (4.16)\end{aligned}$$

The complex variables associated with the $\vartheta$ functions listed in (4.15, 4.16) are given below:

$$\mathsf{q} = \exp\Big(-\frac{1}{2\varrho^2_{\widehat{\alpha}}}\Big),\ \mathsf{z} = -\frac{1}{2}\Big(\frac{x^2}{4}\tau + i\Big),\ \mathsf{z}^{(\jmath)}(\tau)\Big|_{\jmath\in\{1,2\}} = \frac{1}{2}\Big(\tau x\Big(1-(\jmath-1)\frac{x}{2}\Big) - i\Big). \quad (4.17)$$

Analogous to the above summation procedure based on $\vartheta$ function a closed-form expression may also be obtained for the second moment of the electric field (4.11):

$$\begin{aligned}\langle \mathsf{E}^2\rangle^{(\pm)} \;=\; & \frac{1}{2} + \varrho^2_{\widehat{\alpha}} - \frac{\exp(-\varrho^2_{\widehat{\alpha}}/2)}{\sqrt{2\pi\varrho^2_{\widehat{\alpha}}}}\Big(\Re\,[\widehat{\alpha}^2\exp(-2i\omega t)]\Re[\mathcal{T}] \\ & \mp\Re\,[i\widehat{\alpha}^2\exp(-2i\omega t)]\Im[\mathcal{H}]\Big), \qquad (4.18)\end{aligned}$$

where the respective linear sums of $\vartheta_3$ and $\vartheta_4$ functions are abbreviated as follows:

$$\begin{aligned}\mathcal{T} \;=\; & \Big(1-\frac{\epsilon^2x^2}{\widetilde{\Delta}^2}\Big)\exp\Big(i\tau x\Big(1-\frac{x}{4}\Big)\Big)\vartheta_3(\mathsf{q},\mathfrak{z}) \\ & +\frac{\epsilon^2x^2}{\widetilde{\Delta}^2}\exp\Big(-i\tau\Big(1-x+\frac{x^2}{4}\Big)\Big)\vartheta_3(q,\mathfrak{z}^{(1)}), \qquad (4.19) \\ \mathcal{H} \;=\; & \Big[\Big(1-\frac{\epsilon^2}{2\widetilde{\Delta}^2}\Big(1+2x+\frac{11x^2}{2}\Big)\Big)\exp\Big(i\tau x\Big(1-\frac{x}{4}\Big)\Big) + \frac{\epsilon^2}{\widetilde{\Delta}^2}f(1+4x)\times \\ & \times\exp\Big(i\tau x\Big(1-\frac{3x}{4}\Big)\Big) - \frac{5\epsilon^2}{4\widetilde{\Delta}^2}f^2\exp\Big(i\tau x(1-\frac{5x}{4}\Big)\Big)\Big]\,\vartheta_4(\mathsf{q},\mathfrak{z}) \\ & -\frac{\epsilon^2}{\widetilde{\Delta}^2}x\Big(1+\frac{11x}{4}\Big)\exp\Big(-i\tau\Big(1-x+\frac{x^2}{4}\Big)\Big)\vartheta_4(q,\mathfrak{z}^{(1)}) \\ & +\frac{5\epsilon^2}{2\widetilde{\Delta}^2}xf\exp\Big(-i\tau\Big(1-2x+\frac{3x^2}{4}\Big)\Big)\vartheta_4(q,\mathfrak{z}^{(2)}). \qquad (4.20)\end{aligned}$$

The complex variables of the $\vartheta$ functions quoted in (4.19, 4.20) are listed below:

$$\mathfrak{z} = -\frac{1}{2}\Big(\frac{x^2}{2}\tau + i\Big),\quad \mathfrak{z}^{(\jmath)}(\tau)\Big|_{\jmath\in\{1,2\}} = \frac{1}{2}\Big(\tau x\Big(1-(2\jmath-1)\frac{x}{4}\Big) - i\Big). \qquad (4.21)$$

This completes our demonstration of evaluations of the expectation values of antinormally ordered operators in closed forms comprising of the linear combination of $\vartheta$ functions. The procedure is valid for the moderately strong coupling regime where we retain up to quadratic terms in the Laguerre polynomials. In Figs. [6] and [7] we compare, respectively, the expectation values (4.14, 4.18) with the corresponding sums (4.10, 4.11) in the parametric domain referred to in the context of (3.15). We observe good agreement between two sets of values.



Figure 6: First moment of the electric field operator for the values $\lambda = 0.16\,\omega$ and $\alpha = 1$ derived by using (a) the series (4.10), and (b) the analytical evaluation (4.14).



Figure 7: Second moment of the electric field operator for the values $\lambda = 0.16\,\omega$ and $\alpha = 1$ derived using (a) the series (4.11), and (b) the evaluation (4.18) using theta function.

# V Conclusion

We investigated a qubit-oscillator bipartite system in the framework of adiabatic approximation in the regime of moderately strong coupling and also in the presence of a static bias term in the Hamiltonian. Starting with an entangled quasi-Bell state we obtained the evolution of the reduced density matrices of the qubit and the oscillator. The reduced density matrix of the qubit is time independent in the absence of the bias term in the Hamiltonian. Identifying the series with the Jacobi theta functions we evaluated the density matrix elements in closed form as linear combinations of theta functions. The reduced density matrix of the qubit, in turn, yields the entropy of the system. On the other hand the reduced density matrix of the oscillator provides a way to study the phase space density functions of the oscillator degree of freedom. In particular, we evaluate the $Q$-function and utilize this for obtaining closed-form expressions of the first two moments of the scaled electric field operator as linear combinations of theta functions. The first moment of the electric field in the quasi-Bell states assumes non-zero value only in the presence of the bias parameter. Our closed-form evaluations of various physical quantities are compared with, and found to be good approximations of, their series values in the regime characterized by moderately strong coupling as well as weak bias .

The theta function based summation procedure advanced here may be of utility in developing the theory for still larger qubit-oscillator coupling regime. In these instances we need to include higher order terms in the expansion of Laguerre polynomials appearing in the phase factors of the density matrix elements. It is evident from our derivation that the corresponding series still remains convergent owing to the Gaussian weight factor. A generalization of theta functions depending on more variables may be developed to evaluate such quantities in closed form. Another natural direction in which the present approach may be extended is to study a system with a higher number of qubits coupled to an oscillator. These works will be presented elsewhere.

# Acknowledgement

B.Virgin Jenisha would like to acknowledge the fellowship received from University Grants Commission (India) under the scheme Maulana Azad National Fellowship.

# References

[1] E.T. Jaynes, F.W. Cummings, Proc. IEEE **51**, 89 (1963).

[2] A.D.Armour, M.P. Blencowe, K.C. Schwab, Phys. Rev. Lett. **88**, 148301 (2002).

[3] K. Hennessy, Badolato, M. Winger, D. Gerace, M. Atatire, S. Guide, S. Fält, E.L. Hu, A. Imamoglu, Nature (London) **445**, 896 (2007).

[4] R. Letureq, C. Stampfer, K. Inderbizin, L. Durer, C. Hierold, F. Mariani, M.G. Schultz, F. van Oppen, K. Ensslin, Nature Phys. **5**, 327 (2009).

[5] E.K. Irish, J. Gea-Banacloche, J. Martin, K.C. Schwab, Phys. Rev. B **72**, 195410 (2005).

[6] S. Ashhab, F. Nori, Phys. Rev. A **81**, 042311 (2010).

[7] E.K. Irish, Phys. Rev. Lett. **99**, 173601 (2007).

[8] S. Agarwal, S.M. Hashemi Rafsanjani, J.H. Eberly, Phys. Rev. A**85**, 043815 (2012).

[9] G.E. Andrews, R. Askey, R. Roy, *Special Functions*, Cambridge University Press, Cambridge (1999).

[10] J. Hausinger, M. Grifoni, Phys. Rev. A **82**, 062320 (2010).

[11] J.H. Van Vleck, Phys. Rev. **33**, 467 (1929).

[12] P. Yang, P. Zou, Z.M. Zhang, Phys. Lett. A **376**, 2977 (2012).

[13] E. Solano, G.S. Agarwal, H. Walther, Phys. Rev. Lett. **90**, 027903 (2003).

[14] Wolfgang P. Schleich, *Quantum Optics in Phase Space*, Wiley-VCH, Berlin(2001).